

Probing photon structure in DVCS on a photon target

M. El Beiyad ^{a,b}, S. Friot ^c, B.Pire ^a, L. Szymanowski ^d, S. Wallon ^b

^a *CPhT, École Polytechnique, CNRS, Palaiseau, France*

^b *LPT, Université Paris XI, CNRS, Orsay, France*

^c *IPN, CNRS, Orsay, France*

^d *Soltan Institute for Nuclear Studies, Warsaw, Poland*

Abstract

The factorization of the amplitude for the deeply virtual Compton scattering (DVCS) process $\gamma^*(Q)\gamma \rightarrow \gamma\gamma$ at high Q^2 is demonstrated in two distinct kinematical domains, allowing to define the photon generalized parton distributions and the diphoton generalized distribution amplitudes. Both these quantities exhibit an anomalous scaling behaviour and obey new inhomogeneous QCD evolution equations.

Key words: QCD, Exclusive processes, Factorization, Generalized Parton Distributions
PACS: 12.38.Bx, 12.20.Ds, 14.70.Bh

1. Motivation

The parton content of the photon has been the subject of many studies since the seminal paper by Witten [1] which allowed to define the anomalous quark and gluon distribution functions. Recent progresses in exclusive hard reactions focus on generalized parton distributions (GPDs), which are defined as Fourier transforms of matrix elements between different states, such as $\langle N'(p', s') | \bar{\psi}(-\lambda n) \gamma \cdot n \psi(\lambda n) | N(p, s) \rangle$ and their crossed versions, the generalized distribution amplitudes (GDAs) which describe the exclusive hadronization of a $\bar{q}q$ or gg pair in a pair of hadrons, see Fig. 1. In the photon case, these quantities are perturbatively calculable [2,3] at leading order in α_{em} and leading logarithmic order in Q^2 . They constitute an interesting theoretical laboratory for the non-perturbative hadronic objects that hadronic GPDs and GDAs are.

2. The diphoton generalized distribution amplitudes

Defining the momenta as $q = p - \frac{Q^2 n}{s}$, $q' = \frac{Q^2 n}{s}$, $p_1 = \zeta p$, $p_2 = (1 - \zeta)p$, where p and n are two light-cone Sudakov vectors and $2p \cdot n = s$, the amplitude of the process




Fig. 1. Factorizations of the DVCS process on the photon.

$$\gamma^*(Q, \epsilon) \gamma(q', \epsilon') \rightarrow \gamma(p_1, \epsilon_1) \gamma(p_2, \epsilon_2) \quad (1)$$

may be written as $A = \epsilon_\mu \epsilon'_\nu \epsilon_{1\alpha}^* \epsilon_{2\beta}^* T^{\mu\nu\alpha\beta}$. In forward kinematics where $(q + q')^2 = 0$, the tensorial decomposition of $T^{\mu\nu\alpha\beta}$ reads (see [3])

$$\frac{1}{4} g_T^{\mu\nu} g_T^{\alpha\beta} W_1^q + \frac{1}{8} \left(g_T^{\mu\alpha} g_T^{\nu\beta} + g_T^{\nu\alpha} g_T^{\mu\beta} - g_T^{\mu\nu} g_T^{\alpha\beta} \right) W_2^q + \frac{1}{4} \left(g_T^{\mu\alpha} g_T^{\nu\beta} - g_T^{\mu\beta} g_T^{\alpha\nu} \right) W_3^q.$$

At leading order, the three scalar functions W_i^q can be written in a factorized form which is particularly simple when the factorization scale M_F equals the photon virtuality Q . W_1^q is then the convolution $W_1^q = \int_0^1 dz C_V^q(z) \Phi_1^q(z, \zeta, 0)$ of the coefficient function $C_V^q = e_q^2 \left(\frac{1}{z} - \frac{1}{1-z} \right)$ with the *anomalous* vector GDA ($\bar{z} = 1 - z, \bar{\zeta} = 1 - \zeta$) :

$$\Phi_1^q(z, \zeta, 0) = \frac{N_C e_q^2}{2\pi^2} \log \frac{Q^2}{m^2} \left[\frac{\bar{z}(2z - \zeta)}{\bar{\zeta}} \theta(z - \zeta) + \frac{\bar{z}(2z - \bar{\zeta})}{\zeta} \theta(z - \bar{\zeta}) + \frac{z(2z - 1 - \zeta)}{\zeta} \theta(\zeta - z) + \frac{z(2z - 1 - \bar{\zeta})}{\bar{\zeta}} \theta(\bar{\zeta} - z) \right].$$

Conversely, W_3^q is the convolution of the function $C_A^q = e_q^2 \left(\frac{1}{z} + \frac{1}{\bar{z}} \right)$ with the axial GDA :

$$\Phi_3^q(z, \zeta, 0) = \frac{N_C e_q^2}{2\pi^2} \log \frac{Q^2}{m^2} \left[\frac{\bar{z}\zeta}{\bar{\zeta}} \theta(z - \zeta) - \frac{\bar{z}\bar{\zeta}}{\zeta} \theta(z - \bar{\zeta}) - \frac{z\bar{\zeta}}{\zeta} \theta(\zeta - z) + \frac{z\zeta}{\bar{\zeta}} \theta(\bar{\zeta} - z) \right]$$

and $W_2^q = 0$. Note that these GDAs are not continuous at the points $z = \pm\zeta$. The anomalous nature of Φ_1^q and Φ_3^q comes from their proportionality to $\log \frac{Q^2}{m^2}$, which reminds us of the anomalous photon structure functions. A consequence is that $\frac{d}{d \ln Q^2} \Phi_i^q \neq 0$; consequently the QCD evolution equations of the diphoton GDAs obtained with the help of the ERBL kernel are non-homogeneous ones.

3. The photon generalized parton distributions

We now look at the same process in different kinematics, namely $q = -2\xi p + n$, $q' = (1 + \xi)p$, $p_1 = n$, $p_2 = p_1 + \Delta = (1 - \xi)p$, where $W^2 = \frac{1-\xi}{2\xi}Q^2$ and $t = 0$. The tensor $T^{\mu\nu\alpha\beta}$ is now decomposed on different tensors with the help of three functions $\mathcal{W}_i^q$ as (see [2]):

$$\frac{1}{4}g_T^{\mu\alpha}g_T^{\nu\beta}\mathcal{W}_1^q + \frac{1}{8}\left(g_T^{\mu\nu}g_T^{\alpha\beta} + g_T^{\alpha\nu}g_T^{\mu\beta} - g_T^{\mu\alpha}g_T^{\nu\beta}\right)\mathcal{W}_2^q + \frac{1}{4}\left(g_T^{\mu\nu}g_T^{\alpha\beta} - g_T^{\mu\beta}g_T^{\nu\alpha}\right)\mathcal{W}_3^q$$

These functions can also be written in factorized forms which have direct parton model interpretations when the factorization scale M_F is equal to Q . The coefficient functions are $C_{V/A}^q = -2e_q^2\left(\frac{1}{x-\xi+i\eta} \pm \frac{1}{x+\xi-i\eta}\right)$ and the unpolarized H_1^q and polarized H_3^q anomalous GPDs of quarks inside a real photon read :

$$H_1^q(x, \xi, 0) = \frac{N_C e_q^2}{4\pi^2} \left[\theta(x - \xi) \frac{x^2 + (1 - x)^2 - \xi^2}{1 - \xi^2} + \theta(\xi - x)\theta(\xi + x) \frac{x(1 - \xi)}{\xi(1 + \xi)} - \theta(-x - \xi) \frac{x^2 + (1 + x)^2 - \xi^2}{1 - \xi^2} \right] \ln \frac{Q^2}{m^2},$$

$$H_3^q(x, \xi, 0) = \frac{N_C e_q^2}{4\pi^2} \left[\theta(x - \xi) \frac{x^2 - (1 - x)^2 - \xi^2}{1 - \xi^2} - \theta(\xi - x)\theta(\xi + x) \frac{1 - \xi}{1 + \xi} + \theta(-x - \xi) \frac{x^2 - (1 + x)^2 - \xi^2}{1 - \xi^2} \right] \ln \frac{Q^2}{m^2}.$$

Similarly as in the GDA case, the anomalous generalized parton distributions H_i^q are proportional to $\ln \frac{Q^2}{m^2}$. Consequently, the anomalous terms H_i^q supply to the usual homogeneous DGLAP-ERBL evolution equations of GPDs a non-homogeneous term which changes them into non-homogeneous evolution equations.

We do not anticipate a rich phenomenology of these photon GPDs, but in the case of a high luminosity electron - photon collider which is not realistic in the near future. However, the fact that one gets explicit expressions for these GPDs may help to understand the meaning of general theorems such as the polynomiality and positivity [4] constrains or the analyticity structure [5]. For instance, one sees that a D-term is needed when expressing the photon GPDs in terms of a double distribution. One also finds that, in the DGLAP region, $H_1(x, \xi)$ is smaller than its positivity bound by a sizeable and slowly varying factor, which is of the order of $0.7 - 0.8$ for $\xi \approx 0.3$.

This work is supported by the Polish Grant N202 249235, the French-Polish scientific agreement Polonium and the grant ANR-06-JCJC-0084.

References

- [1] E. Witten, Nucl. Phys. B **120** (1977) 189.
- [2] S. Friot *et.al.*, Phys. Lett. B **645** (2007) 153 and Nucl. Phys. Proc. Suppl. **184**, 35 (2008).
- [3] M. El Beiyad *et.al.*, Phys. Rev. D **78** (2008) 034009 and AIP Conf. Proc. **1038**, 305 (2008).

- [4] B. Pire, J. Soffer and O. Teryaev, Eur. Phys. J. C **8** (1999) 103.
- [5] I. V. Anikin *et.al.*, Phys. Rev. D **76**, 056007 (2007). M. Diehl *et.al.*, Eur. Phys. J. C **52** (2007) 919.